


## Lesson 7: Secant and the Co-Functions

### Classwork

#### Opening Exercise

Find the length of each segment below in terms of the value of a trigonometric function.

$OQ =$  \_\_\_\_\_       $PQ =$  \_\_\_\_\_       $RS =$  \_\_\_\_\_



#### Example 1

Use similar triangles to find the value of  $\sec(\theta^\circ)$  in terms of one other trigonometric function.

**Exercise 1**

The definition of the secant function is offered below. Answer the questions to better understand this definition and the domain and range of this function. Be prepared to discuss your responses with others in your class.


Let  $\theta$  be any real number.

In the Cartesian plane, rotate the nonnegative  $x$ -axis by  $\theta$  degrees about the origin. Intersect this new ray with the unit circle to get a point  $(x_\theta, y_\theta)$ .

If  $x_\theta \neq 0$ , then the value of  $\sec(\theta^\circ)$  is  $\frac{1}{x_\theta}$ .

Otherwise,  $\sec(\theta^\circ)$  is undefined.


In terms of the cosine function,  $\sec(\theta^\circ) = \frac{1}{\cos(\theta^\circ)}$  for  $\cos(\theta^\circ) \neq 0$ .



- a. What is the domain of the secant function?
  
- b. The domains of the secant and tangent functions are the same. Why?
  
- c. What is the range of the secant function? How is this range related to the range of the cosine function?
  
- d. Is the secant function a periodic function? If so, what is its period?


**Exercise 2**

In the diagram, the horizontal blue line is tangent to the unit circle at  $(0,1)$ .



- How does this diagram compare to the one given in the Opening Exercise?
- What is the relationship between  $\beta$  and  $\theta$ ?
- Which segment in the figure has length  $\sin(\theta^\circ)$ ? Which segment has length  $\cos(\theta^\circ)$ ?
- Which segment in the figure has length  $\sin(\beta^\circ)$ ? Which segment has length  $\cos(\beta^\circ)$ ?
- How can you write  $\sin(\theta^\circ)$  and  $\cos(\theta^\circ)$  in terms of the trigonometric functions of  $\beta$ ?

Example 2



The horizontal blue line is tangent to the circle at  $(0,1)$ .

- a. If two angles are complements with measures  $\beta$  and  $\theta$  as shown in the diagram, use similar triangles to show that  $\sec(\beta^\circ) = \frac{1}{\sin(\theta^\circ)}$ .

- b. If two angles are complements with measures  $\beta$  and  $\theta$  as shown in the diagram, use similar triangles to show that  $\tan(\beta^\circ) = \frac{1}{\tan(\theta^\circ)}$ .

**Discussion**


Definitions of the cosecant and cotangent functions are offered below. Answer the questions to better understand the definitions and the domains and ranges of these functions. Be prepared to discuss your responses with others in your class.

Let  $\theta$  be any real number such that  $\theta \neq 180k$  for all integers  $k$ .

In the Cartesian plane, rotate the initial ray by  $\theta$  degrees about the origin. Intersect the resulting terminal ray with the unit circle to get a point  $(x_\theta, y_\theta)$ .

The value of  $\csc(\theta^\circ)$  is  $\frac{1}{y_\theta}$ .

The value of  $\cot(\theta^\circ)$  is  $\frac{x_\theta}{y_\theta}$ .



The secant, cosecant, and cotangent functions are often referred to as reciprocal functions. Why do you think these functions are so named?

Why are the domains of these functions restricted?

The domains of the cosecant and cotangent functions are the same. Why?

What is the range of the cosecant function? How is this range related to the range of the sine function?


What is the range of the cotangent function? How is this range related to the range of the tangent function?

Let  $\theta$  be any real number. In the Cartesian plane, rotate the initial ray by  $\theta$  degrees about the origin. Intersect the resulting terminal ray with the unit circle to get a point  $(x_\theta, y_\theta)$ . Then:

Function	Value	For any $\theta$ such that...	Formula
Sine	$y_\theta$	$\theta$ is a real number	
Cosine	$x_\theta$	$\theta$ is a real number	
Tangent	$\frac{y_\theta}{x_\theta}$	$\theta \neq 90 + 180k$ , for all integers $k$	$\tan(\theta^\circ) = \frac{\sin(\theta^\circ)}{\cos(\theta^\circ)}$
Secant	$\frac{1}{x_\theta}$	$\theta \neq 90 + 180k$ , for all integers $k$	$\sec(\theta^\circ) = \frac{1}{\cos(\theta^\circ)}$
Cosecant	$\frac{1}{y_\theta}$	$\theta \neq 180k$ , for all integers $k$	$\csc(\theta^\circ) = \frac{1}{\sin(\theta^\circ)}$
Cotangent	$\frac{x_\theta}{y_\theta}$	$\theta \neq 180k$ , for all integers $k$	$\cot(\theta^\circ) = \frac{\cos(\theta^\circ)}{\sin(\theta^\circ)}$

**Problem Set**

- Use the reciprocal interpretations of  $\sec(\theta^\circ)$ ,  $\csc(\theta^\circ)$ , and  $\cot(\theta^\circ)$  and the unit circle provided to complete the table.



$\theta$ , in degrees	$\sec(\theta^\circ)$	$\csc(\theta^\circ)$	$\cot(\theta^\circ)$
0			
30			
45			
60			
90			
120			
180			
225			
240			
270			
315			
330			

2. Find the following values from the information given.
- $\sec(\theta^\circ)$ ;  $\cos(\theta^\circ) = 0.3$
  - $\csc(\theta^\circ)$ ;  $\sin(\theta^\circ) = -0.05$
  - $\cot(\theta^\circ)$ ;  $\tan(\theta^\circ) = 1000$
  - $\sec(\theta^\circ)$ ;  $\cos(\theta^\circ) = -0.9$
  - $\csc(\theta^\circ)$ ;  $\sin(\theta^\circ) = 0$
  - $\cot(\theta^\circ)$ ;  $\tan(\theta^\circ) = -0.0005$
3. Choose three  $\theta$  values from the table in Problem 1 for which  $\sec(\theta^\circ)$ ,  $\csc(\theta^\circ)$ , and  $\tan(\theta^\circ)$  are defined and not zero. Show that for these values of  $\theta$ ,  $\frac{\sec(\theta^\circ)}{\csc(\theta^\circ)} = \tan(\theta^\circ)$ .
4. Find the value of  $\sec(\theta^\circ)\cos(\theta^\circ)$  for the following values of  $\theta$ .
- $\theta = 120$
  - $\theta = 225$
  - $\theta = 330$
  - Explain the reasons for the pattern you see in your responses to parts (a)–(c).
5. Draw a diagram representing the two values of  $\theta$  between 0 and 360 so that  $\sin(\theta^\circ) = -\frac{\sqrt{3}}{2}$ . Find the values of  $\tan(\theta^\circ)$ ,  $\sec(\theta^\circ)$ , and  $\csc(\theta^\circ)$  for each value of  $\theta$ .
6. Find the value of  $(\sec(\theta^\circ))^2 - (\tan(\theta^\circ))^2$  when  $\theta = 225$ .
7. Find the value of  $(\csc(\theta^\circ))^2 - (\cot(\theta^\circ))^2$  when  $\theta = 330$ .

**Extension:**

8. Using the formulas  $\sec(\theta^\circ) = \frac{1}{\cos(\theta^\circ)}$ ,  $\csc(\theta^\circ) = \frac{1}{\sin(\theta^\circ)}$ , and  $\cot(\theta^\circ) = \frac{1}{\tan(\theta^\circ)}$ , show that  $\frac{\sec(\theta^\circ)}{\csc(\theta^\circ)} = \tan(\theta^\circ)$ , where these functions are defined and not zero.
9. Tara showed that  $\frac{\sec(\theta^\circ)}{\csc(\theta^\circ)} = \tan(\theta^\circ)$ , for values of  $\theta$  for which the functions are defined and  $\csc(\theta^\circ) \neq 0$ , and then concluded that  $\sec(\theta^\circ) = \sin(\theta^\circ)$  and  $\csc(\theta^\circ) = \cos(\theta^\circ)$ . Explain what is wrong with her reasoning.
10. From Lesson 6, Ren remembered that the tangent function is odd, meaning that  $-\tan(\theta^\circ) = \tan(-\theta^\circ)$  for all  $\theta$  in the domain of the tangent function. He concluded because of the relationship between the secant function, cosecant function, and tangent function developed in Problem 9, it is impossible for both the secant and the cosecant functions to be odd. Explain why he is correct.