

Lesson 4: Properties of Exponents and Radicals

Classwork

Opening Exercise

Write each exponential expression as a radical expression, and then use the definition and properties of radicals to write the resulting expression as an integer.

a. $7^{\frac{1}{2}} \cdot 7^{\frac{1}{2}}$

b. $3^{\frac{1}{3}} \cdot 3^{\frac{1}{3}} \cdot 3^{\frac{1}{3}}$

c. $12^{\frac{1}{2}} \cdot 3^{\frac{1}{2}}$

d. $\left(64^{\frac{1}{3}}\right)^{\frac{1}{2}}$

Examples 1–3

Write each expression in the form $b^{\frac{m}{n}}$ for positive real numbers b and integers m and n with $n > 0$ by applying the properties of radicals and the definition of n^{th} root.

1. $b^{\frac{1}{4}} \cdot b^{\frac{1}{4}}$

2. $b^{\frac{1}{3}} \cdot b^{\frac{4}{3}}$

3. $b^{\frac{1}{5}} \cdot b^{\frac{3}{4}}$

Exercises 1–4

Write each expression in the form $b^{\frac{m}{n}}$. If a numeric expression is a rational number, then write your answer without exponents.

1. $b^{\frac{2}{3}} \cdot b^{\frac{1}{2}}$

2. $\left(b^{-\frac{1}{5}}\right)^{\frac{2}{3}}$

3. $64^{\frac{1}{3}} \cdot 64^{\frac{3}{2}}$

4. $\left(\frac{9^3}{4^2}\right)^{\frac{3}{2}}$

Example 4

Rewrite the radical expression $\sqrt{48x^5y^4z^2}$ so that no perfect square factors remain inside the radical.

Exercise 5

5. Use the definition of rational exponents and properties of exponents to rewrite each expression with rational exponents containing as few fractions as possible. Then, evaluate each resulting expression for $x = 50$, $y = 12$, and $z = 3$.

a. $\sqrt{8x^3y^2}$

b. $\sqrt[3]{54y^7z^2}$

Exercise 6

6. Order these numbers from smallest to largest. Explain your reasoning.

$16^{2.5}$

$9^{3.6}$

$32^{1.2}$

Lesson Summary

The properties of exponents developed in Grade 8 for integer exponents extend to rational exponents.

That is, for any integers $m, n, p,$ and $q,$ with $n > 0$ and $q > 0,$ and any real numbers a and b so that $a^{\frac{1}{n}}, b^{\frac{1}{n}},$ and $b^{\frac{1}{q}}$ are defined, we have the following properties of exponents:

1. $b^{\frac{m}{n}} \cdot b^{\frac{p}{q}} = b^{\frac{m+p}{q}}$
2. $b^{\frac{m}{n}} = \sqrt[n]{b^m}$
3. $\left(b^{\frac{1}{n}}\right)^n = b$
4. $\left(b^n\right)^{\frac{1}{n}} = b$
5. $(ab)^{\frac{m}{n}} = a^{\frac{m}{n}} \cdot b^{\frac{m}{n}}$
6. $\left(b^{\frac{m}{n}}\right)^{\frac{p}{q}} = b^{\frac{mp}{nq}}$
7. $b^{-\frac{m}{n}} = \frac{1}{b^{\frac{m}{n}}}$

Problem Set

1. Evaluate each expression for $a = 27$ and $b = 64.$

- | | |
|---|--|
| a. $\sqrt[3]{a}\sqrt{b}$ | d. $a^{-\frac{2}{3}} + b^{\frac{3}{2}}$ |
| b. $\left(3\sqrt[3]{a}\sqrt{b}\right)^2$ | e. $\left(a^{-\frac{2}{3}} \cdot b^{\frac{3}{2}}\right)^{-1}$ |
| c. $\left(\sqrt[3]{a} + 2\sqrt{b}\right)^2$ | f. $\left(a^{-\frac{2}{3}} - \frac{1}{8}b^{\frac{3}{2}}\right)^{-1}$ |

2. Rewrite each expression so that each term is in the form $kx^n,$ where k is a real number, x is a positive real number, and n is a rational number.

- | | |
|---|--|
| a. $x^{-\frac{2}{3}} \cdot x^{\frac{1}{3}}$ | f. $\sqrt[3]{\frac{27}{x^6}}$ |
| b. $2x^{\frac{1}{2}} \cdot 4x^{-\frac{5}{2}}$ | g. $\sqrt[3]{x} \cdot \sqrt[3]{-8x^2} \cdot \sqrt[3]{27x^4}$ |
| c. $\frac{10x^{\frac{1}{3}}}{2x^2}$ | h. $\frac{2x^4 - x^2 - 3x}{\sqrt{x}}$ |
| d. $\left(3x^{\frac{1}{4}}\right)^{-2}$ | i. $\frac{\sqrt{x} - 2x^{-3}}{4x^2}$ |
| e. $x^{\frac{1}{2}}\left(2x^2 - \frac{4}{x}\right)$ | |

3. Show that $(\sqrt{x} + \sqrt{y})^2$ is not equal to $x^1 + y^1$ when $x = 9$ and $y = 16$.
4. Show that $(x^{\frac{1}{2}} + y^{\frac{1}{2}})^{-1}$ is not equal to $\frac{1}{x^{\frac{1}{2}}} + \frac{1}{y^{\frac{1}{2}}}$ when $x = 9$ and $y = 16$.
5. From these numbers, select (a) one that is negative, (b) one that is irrational, (c) one that is not a real number, and (d) one that is a perfect square:

$$\frac{1}{3^2} \cdot \frac{1}{9^2}, 27^{\frac{1}{3}} \cdot 144^{\frac{1}{2}}, 64^{\frac{1}{3}} - 64^{\frac{2}{3}}, \text{ and } \left(4^{-\frac{1}{2}} - 4^{\frac{1}{2}}\right)^{\frac{1}{2}}$$

6. Show that for any rational number n , the expression $2^n \cdot 4^{n+1} \cdot \left(\frac{1}{8}\right)^n$ is equal to 4.
7. Let n be any rational number. Express each answer as a power of 10.
- Multiply 10^n by 10.
 - Multiply $\sqrt{10}$ by 10^n .
 - Square 10^n .
 - Divide $100 \cdot 10^n$ by 10^{2n} .
 - Show that $10^n = 11 \cdot 10^n - 10^{n+1}$.
8. Rewrite each of the following radical expressions as an equivalent exponential expression in which each variable occurs no more than once.
- $\sqrt{8x^2y}$
 - $\sqrt[5]{96x^3y^{15}z^6}$
9. Use properties of exponents to find two integers that are upper and lower estimates of the value of $4^{1.6}$.
10. Use properties of exponents to find two integers that are upper and lower estimates of the value of $8^{2.3}$.
11. Kepler's third law of planetary motion relates the average distance, a , of a planet from the sun to the time, t , it takes the planet to complete one full orbit around the sun according to the equation $t^2 = a^3$. When the time, t , is measured in Earth years, the distance, a , is measured in astronomical units (AUs). (One AU is equal to the average distance from Earth to the sun.)
- Find an equation for t in terms of a and an equation for a in terms of t .
 - Venus takes about 0.616 Earth year to orbit the sun. What is its average distance from the sun?
 - Mercury is an average distance of 0.387 AU from the sun. About how long is its orbit in Earth years?