


Lesson 8: How Do Dilations Map Lines, Rays, and Circles?

Classwork

Opening Exercise

- a. Is a dilated ray still a ray? If the ray is transformed under a dilation, explain how.
- b. Dilate the \overrightarrow{PQ} by a scale factor of 2 from center O .



- i. Is the figure $\overrightarrow{P'Q'}$ a ray?

- ii. How, if at all, has the ray PQ been transformed?
- iii. Will a ray always be mapped to a ray? Explain how you know.

Example 1

Will a dilation about center O and scale factor $r = 1$ map \overrightarrow{PQ} to $\overrightarrow{P'Q'}$? Explain.

Example 2

The line that contains \overrightarrow{PQ} does not contain point O . Does a dilation about center O and scale factor $r \neq 1$ map every point of \overrightarrow{PQ} onto a point of $\overrightarrow{P'Q'}$?

Example 3

The line that contains \overrightarrow{PQ} contains point O . Does a dilation about center O and scale factor r map \overrightarrow{PQ} to $\overrightarrow{P'Q'}$?

- Examine the case where the endpoint P of \overrightarrow{PQ} coincides with the center O of the dilation.

- Examine the case where the endpoint P of \overrightarrow{PQ} is between O and Q on the line containing O , P , and Q .

- Examine the remaining case where the center O of the dilation and point Q are on the same side of P on the line containing O , P , and Q .

Example 5

Does a dilation about a center O and scale factor r map a circle of radius R onto another circle?

- Examine the case where the center of the dilation coincides with the center of the circle.

- Examine the case where the center of the dilation is not the center of the circle; we call this the *general case*.

Lesson Summary

- **DILATION THEOREM FOR RAYS:** A dilation maps a ray to a ray sending the endpoint to the endpoint.
- **DILATION THEOREM FOR LINES:** A dilation maps a line to a line. If the center O of the dilation lies on the line or if the scale factor r of the dilation is equal to 1, then the dilation maps the line to the same line. Otherwise, the dilation maps the line to a parallel line.
- **DILATION THEOREM FOR CIRCLES:** A dilation maps a circle to a circle and maps the center to the center.

Problem Set


1. In Lesson 8, Example 2, you proved that a dilation with a scale factor $r > 1$ maps a ray PQ to a ray $P'Q'$. Prove the remaining case that a dilation with scale factor $0 < r < 1$ maps a ray PQ to a ray $P'Q'$.

Given the dilation $D_{O,r}$, with $0 < r < 1$ maps P to P' and Q to Q' , prove that $D_{O,r}$ maps \overrightarrow{PQ} to $\overrightarrow{P'Q'}$.

2. In the diagram below, $\overrightarrow{A'B'}$ is the image of \overrightarrow{AB} under a dilation from point O with an unknown scale factor; A maps to A' , and B maps to B' . Use direct measurement to determine the scale factor r , and then find the center of dilation O .



3. Draw a line \overleftrightarrow{AB} , and dilate points A and B from center O where O is not on \overleftrightarrow{AB} . Use your diagram to explain why a line maps to a line under a dilation with scale factor r .
4. Let \overline{AB} be a line segment, and let m be a line that is the perpendicular bisector of \overline{AB} . If a dilation with scale factor r maps \overline{AB} to $\overline{A'B'}$ (sending A to A' and B to B') and also maps line m to line m' , show that line m' is the perpendicular bisector of $\overline{A'B'}$.
5. Dilate circle C with radius CA from center O with a scale factor $r = \frac{1}{2}$.



6. In the picture below, the larger circle is a dilation of the smaller circle. Find the center of dilation O .

